

Long-time behaviors and stability of entropy solutions for linearly degenerate hyperbolic systems of rich type

Yue-Jun PENG^{1,2} and Yong-Fu YANG^{3,4}

¹Clermont Université, Université Blaise Pascal, 63000 Clermont-Ferrand, France

²CNRS-UMR 6620, Laboratoire de Mathématiques, 63171 Aubière cedex, France

³LCP, Institute of Applied Physics and Computational Mathematics, Beijing, 100088, China

⁴Department of Mathematics, College of Sciences, Hohai University, Nanjing 210098, China

Email : peng@math.univ-bpclermont.fr, yongfuyang@sina.com

Abstract. We show that in one space dimension, a linearly degenerate hyperbolic system of rich type admits exact traveling wave solutions if the initial data are Riemann type outside of a space interval. In a particular case of the system including physical models, we prove the convergence of entropy solutions to traveling waves in the L^1 norm as the time goes to infinity. The traveling waves are determined explicitly in terms of the initial data and the system. We also obtain the stability of entropy solutions in L^1 .

Keywords. Entropy solution, linearly degeneracy, rich system, non-strict hyperbolicity, long-time behavior, traveling wave, L^1 stability

1 Introduction

For first order quasilinear hyperbolic systems of conservation laws, it is well known that, generically speaking, classical solutions exist only locally in time and singularities may appear in a finite time. Therefore, we are often led to seek global entropy solutions (see [10, 14, 25] and the references therein). In some special cases, however, the global existence of classical solutions can be obtained. In particular, these situations occur for linearly degenerate systems provided that the initial data are sufficiently small and decay at infinity (see [6, 15, 17, 18]).

As to the Cauchy problem in one space dimension, the existence of global entropy solutions was proved by Glimm [8] when the initial data are small in the total variation norm. The long-time behavior of the entropy solution was investigated by Glimm and Lax [9] for systems of two conservation laws having genuinely nonlinear characteristics. In [20] Liu studied this problem for a general hyperbolic system of conservation laws when the initial data agree with Riemann data outside of a large space interval. He showed that waves of genuinely nonlinear characteristic fields tend to the solution of the corresponding Riemann problem and waves of linearly degenerate characteristic fields tend to traveling waves, both at algebraic rates. The

convergence is in the BV norm. When the initial data have compact support, the solution also converges in the L^1 norm to the linear superposition of traveling waves and N -waves.

If the system is strictly hyperbolic and weakly linearly degenerate (see [15] for definition), Kong and Yang [13] also discussed the long-time behavior of global classical solutions to its Cauchy problem with small and decaying initial data, provided that the source term satisfies a so-called matching condition. Recently, Liu and Zhou [19] considered the similar long-time behavior of classical solutions for diagonal systems when the initial data are large. They applied their results to the generalized extremal surface equations. All these results indicate that the classical solutions of a linearly degenerate hyperbolic system approach some C^1 traveling wave solutions. Unfortunately, all these results are qualitative on the limits, from which we do not know what these traveling waves are.

Stability is another important problem for hyperbolic systems of conservation laws. In particular, it implies the uniqueness of solutions. A complete theory of the uniqueness and L^1 stability of entropy solutions to the scalar conservation law was established by Kruzkov [11]. See also [1] and the references therein by a kinetic method. For hyperbolic systems of conservation laws, we refer to the work of Bressan [4] for the existence, uniqueness and stability of entropy solutions in $L^\infty(\mathbb{R}) \cap BV(\mathbb{R})$ with small initial data.

In a previous work, the authors of this paper considered the generalized extremal surface equations (see [2, 12]) and showed that its Lipschitz solutions are equivalent to entropy solutions in $L^\infty(\mathbb{R})$ of a non-strictly hyperbolic system of conservation laws [23]. They obtained an explicit representation formula and the uniqueness of the entropy solutions to the Cauchy problem. Based on these results, they further obtained the convergence and convergence rates of the entropy solutions to traveling waves in the $L^1(\mathbb{R})$ norm. The traveling waves are determined explicitly in terms of the initial data and the system. Moreover, when initial data agree with Riemann data outside of a finite space interval, the entropy solutions become explicit traveling waves after a finite time. Finally, they also got L^1 stability of the entropy solutions.

In this paper, we are going to extend results of [23] to linearly degenerate hyperbolic systems of rich type. The notion of rich system was introduced by Serre [24] and Tsarëv [28, 29] independently. It is also called semi-hamiltonian system by Tsarëv. For this kind of linearly degenerate system, the global existence of entropy solutions in L^∞ was established by Chen [7] in the strictly hyperbolic case and in [16] in the non-strictly hyperbolic case. The uniqueness of entropy solutions has not been solved yet. However, it is solved in a particular case where $N_i = N$ (see the definition below) for all i . This case contains many interesting physical models, such as the generalized extremal surface equations, the Born-Infeld system and augmented Born-Infeld system. All these models are non-strictly hyperbolic systems.

Our study on the long-time behavior and stability of entropy solutions in L^1 is carried out in this special case. The main results are stated in Theorems 3.1 and 4.1, which are available for initial data in $L^1 \cap L^\infty$ and also in C^k with integer $k \geq 1$. The traveling wave solution after a finite time holds in a more general case, provided the initial data are Riemann type outside of a finite space interval (see Theorem 2.1). Below are preliminary descriptions on the rich system and related results.

Consider the Cauchy problem for quasilinear hyperbolic systems of diagonal form :

$$\partial_t w_i + \lambda_i(w) \partial_x w_i = 0, \quad \forall i \in K_n, \quad t > 0, \quad x \in \mathbb{R} \quad (1.1)$$

with initial data

$$t = 0 : \quad w = w^0(x), \quad x \in \mathbb{R}. \quad (1.2)$$

Here $K_n = \{1, \dots, n\}$, $w = (w_1, \dots, w_n)^\top$ and $w^0 = (w_1^0, \dots, w_n^0)^\top$, in which $\top$ means transpose. The eigenvalues $\lambda_i(w)$ ($i \in K_n$) of the system are real valued smooth functions defined

on an open domain of $\mathbb{R}^n$. We suppose that each of them has a constant multiplicity, namely, on the domain under consideration we have

$$\begin{aligned} \mu_1(w) \stackrel{\text{def}}{=} \lambda_1(w) = \cdots = \lambda_{r_1}(w) < \mu_2(w) \stackrel{\text{def}}{=} \lambda_{r_1+1}(w) = \cdots = \lambda_{r_2}(w) \\ < \cdots < \mu_s(w) \stackrel{\text{def}}{=} \lambda_{r_{s-1}+1}(w) = \cdots = \lambda_{r_s}(w), \end{aligned} \quad (1.3)$$

where $r_1, r_2, \dots, r_s$ are constants and

$$1 \leq r_1 < r_2 < \cdots < r_s = n.$$

For system (1.1) and $p \in K_s$, the characteristic $\mu_p(w)$ with constant multiplicity $r_p - r_{p-1}$ ($r_0 = 0$) is linearly degenerate if and only if $\mu_p(w)$ is independent of $w_{r_{p-1}+1}, \dots, w_{r_p}$:

$$\frac{\partial \mu_p(w)}{\partial w_l} \equiv 0, \quad (l = r_{p-1} + 1, \dots, r_p).$$

If all characteristics $\lambda_i(w)$ ($i \in K_n$) are linearly degenerate, system (1.1) is said to be linearly degenerate. Let $\Lambda(w) = \text{diag}\{\lambda_1(w), \dots, \lambda_n(w)\}$. A pair of functions $(E(w), F(w))$ is an entropy-entropy flux pair of (1.1) if $F'(w) = E'(w)\Lambda(w)$, i.e.,

$$\frac{\partial F(w)}{\partial w_j} = \lambda_j(w) \frac{\partial E(w)}{\partial w_j}, \quad \forall j \in K_n.$$

It is well known that $(E(w), F(w))$ is an entropy-entropy flux pair if and only if it satisfies the additional conservation law (see [14, 25])

$$\partial_t E(w) + \partial_x F(w) = 0.$$

Recall that the i th characteristic is rich if, on the domain under consideration, for all $j, k \in K_n$ such that $\lambda_j(w) \neq \lambda_i(w)$ and $\lambda_k(w) \neq \lambda_i(w)$, we have

$$\frac{\partial}{\partial w_j} \left(\frac{\frac{\partial \lambda_i(w)}{\partial w_k}}{\lambda_k(w) - \lambda_i(w)} \right) = \frac{\partial}{\partial w_k} \left(\frac{\frac{\partial \lambda_i(w)}{\partial w_j}}{\lambda_j(w) - \lambda_i(w)} \right).$$

This definition is equivalent to the existence of a smooth function $N_i(w) > 0$ such that

$$(\lambda_j(w) - \lambda_i(w)) \frac{\partial N_i(w)}{\partial w_j} = N_i(w) \frac{\lambda_i(w)}{\partial w_j}, \quad \forall j \in K_n, \quad \lambda_j(w) \neq \lambda_i(w). \quad (1.4)$$

When $\lambda_i(w)$ is linearly degenerate, (1.4) is equivalent to

$$(\lambda_j(w) - \lambda_i(w)) \frac{\partial N_i(w)}{\partial w_j} = N_i(w) \frac{\lambda_i(w)}{\partial w_j}, \quad \forall j \in K_n.$$

System (1.1) is said to be rich if all characteristics $\lambda_i(w)$ ($i \in K_n$) are rich. We refer to [25, 27, 29] for the definition and properties of the rich system.

For any given smooth solution $w = w(t, x)$ to the linearly degenerate strictly hyperbolic rich system (1.1), from [25] we have

$$\partial_t N_i(w) + \partial_x (N_i(w) \lambda_i(w)) = 0, \quad i \in K_n. \quad (1.5)$$

Therefore, $(N_i(w), N_i(w) \lambda_i(w))$ is an entropy-entropy flux pair of system (1.1). More generally, we have:

$$\partial_t (N_i(w) g_i(w_i)) + \partial_x (N_i(w) \lambda_i(w) g_i(w_i)) = 0 \quad \text{for any smooth function } g_i. \quad (1.6)$$

Thus, $N_i(w)g_i(w_i)$ ($i \in K_n$) stand for n independent families of entropies and each entropy of the linearly degenerate rich system (1.1) is a linear combination of $N_i(w)g_i(w_i)$ ($i \in K_n$) (see [24, 25]). In particular, taking $g_i(s) = s$ in (1.6) gives :

$$\partial_t (N_i(w)w_i) + \partial_x (N_i(w)\lambda_i(w)w_i) = 0. \quad (1.7)$$

Definition 1.1. A function $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ is an entropy solution to a linearly degenerate hyperbolic rich system of diagonal form (1.1) if the entropy equalities (1.5) and (1.7) are satisfied in the sense of distribution for all $i \in K_n$.

Let $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ be an entropy solution. For each $i \in K_n$, as a compatibility condition, (1.5) implies that there is a function $y_i = Y_i(t, x)$ such that

$$dy_i = N_i(w) dx - (N_i\lambda_i)(w) dt. \quad (1.8)$$

Since

$$\frac{\partial Y_i}{\partial x}|_{t=0} = N_i(w(t, x))|_{t=0} = N_i(w^0(x)),$$

it is natural to define :

$$Y_i(0, x) = Y_i^0(x) \stackrel{\text{def}}{=} \int_0^x N_i(w^0(\xi)) d\xi. \quad (1.9)$$

For any given $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$, (1.8) and (1.9) determine a unique Lipschitz and strictly increasing function $y_i = Y_i(t, x)$ for all $t \geq 0$. As a consequence, from Proposition 2 and Theorem 3 in [16], we have

Proposition 1.1. Suppose that $w^0 \in L^\infty(\mathbb{R})$ and system (1.1) is linearly degenerate and rich. Then the Cauchy problem (1.1)-(1.2) admits a global entropy solution $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ in the sense of Definition 1.1. Moreover,

$$w_i(t, x) = w_i^0(X_i^0(Y_i(t, x))) \quad (i \in K_n), \quad (1.10)$$

where $X_i^0 = (Y_i^0)^{-1}$ is the inverse function of $Y_i^0(\cdot)$. Moreover, if $w^0 \in L^\infty(\mathbb{R}) \cap C^k(\mathbb{R})$ for some integer $k \geq 1$, then $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R}) \cap C^k(\mathbb{R}^+ \times \mathbb{R})$.

Remark 1.1. Proposition 1.1 only establishes the existence of the entropy solutions. Up to our knowledge, so far no uniqueness results are available in this general case of the rich system.

Next we consider the special case that $N_i = N$ for all $i \in K_n$. Then (1.5) becomes :

$$\partial_t N(w) + \partial_x (N(w)\lambda_i(w)) = 0, \quad i \in K_n.$$

From [16], we know that for each $i \in K_n$, there are a constant $\tilde{\lambda}_i$ and a function M independent of i such that

$$N(w)\lambda_i(w) - \tilde{\lambda}_i = M(w). \quad (1.11)$$

Hence,

$$\partial_t N(w) + \partial_x M(w) = 0.$$

This scalar equation allows to make a change of coordinates $(t, x) \mapsto (s, z)$ with (see [30, 22])

$$s = t, \quad dz = N dx - M dt. \quad (1.12)$$

For $w^0 \in L^\infty(\mathbb{R})$, let $z = Z(t, x)$ be the unique Lipschitz function satisfying (1.12) and

$$Z(0, x) = Z^0(x) \stackrel{\text{def}}{=} \int_0^x N(w^0(\xi)) d\xi. \quad (1.13)$$

Since $x \mapsto Z(t, x)$ is a strictly increasing function, we may define its inverse by $X(t, \cdot) = Z^{-1}(t, \cdot)$ for all $t \geq 0$. The function $x = X(t, x)$ satisfies

$$dx = \frac{1}{N(\tilde{w}(t, z))} dz + \frac{M(\tilde{w}(t, z))}{N(\tilde{w}(t, z))} dt, \quad (1.14)$$

where

$$\tilde{w}(t, z) = (\tilde{w}_1^0(z - \tilde{\lambda}_1 t), \dots, \tilde{w}_n^0(z - \tilde{\lambda}_n t))^\top, \quad \tilde{w}_i^0 = w_i^0 \circ X^0. \quad (1.15)$$

Together with $X(0, z) = X^0(z)$, (1.14) determines a unique Lipschitz function for all $t \geq 0$. Then the unique entropy solution of the Cauchy problem (1.1)-(1.2) is given by

$$w_i(t, x) = w_i^0(X^0(Z(t, x) - \tilde{\lambda}_i t)), \quad \forall i \in K_n. \quad (1.16)$$

Moreover, a straightforward computation using (1.11) gives

$$X^0(z) = \int_0^z \frac{1}{N(\tilde{w}^0(z))} dz, \quad X(t, z) = X^0(z) + \int_0^t \left(\lambda_i(\tilde{w}(\tau, z)) - \frac{\tilde{\lambda}_i}{N(\tilde{w}(\tau, z))} \right) d\tau. \quad (1.17)$$

Thus, Li *et al.* [16] got the following result.

Proposition 1.2. *Assume that system (1.1) is linearly degenerate and rich with N_i being independent of i for all $i \in K_n$. Then the Cauchy problem (1.1)-(1.2) with the initial data $w^0 \in L^\infty(\mathbb{R})$ admits a unique entropy solution $w \in L^\infty(\mathbb{R}^+ \times \mathbb{R})$ given by expression (1.16), where X^0 is the inverse function of Z^0 being defined by (1.13) and $Z(t, \cdot)$ is the inverse function of the unique solution $X(t, \cdot)$ to (1.14) and $X(0, z) = X^0(z)$.*

This paper is organized as follows. In the next section, we first derive the explicit traveling waves after a finite time. Section 3 and Section 4 are devoted to the study of the long-time behavior and L^1 stability of the entropy solutions in the case $N_i = N$ for all i , respectively. Finally, we give two examples in Section 5.

2 Explicit traveling waves after a finite time

Throughout this section, we suppose that the initial data w^0 are constants outside of a finite space interval $(-L, L)$ with $L > 0$. More precisely, let

$$w^\pm(x) = \bar{w}^\pm \stackrel{\text{def}}{=} (\bar{w}_1^\pm, \dots, \bar{w}_n^\pm)^\top, \quad \forall \pm x \geq L, \quad (2.1)$$

where $\bar{w}_i^\pm$ are constants for all $i \in K_n$. In addition, to simplify the presentation, we suppose temporarily that system (1.1) is strictly hyperbolic. Based on the explicit expression (1.10), we show that the entropy solution is exactly composed of traveling waves after a finite time $t_L > 0$ to be determined later. In particular, each Riemann invariant is a traveling wave after t_L .

For each $i \in K_n$, since $N_i(w) > 0$, (1.8) implies that $x \mapsto Y_i(t, x)$ is a strictly increasing function and then a bijection for all $t \geq 0$. Let $X_i(t, \cdot)$ be the inverse function of $Y_i(t, \cdot)$ and

$$X_i^\pm(t) = X_i(t, Y_i^0(\pm L)).$$

It follows from (1.10) and (2.1) that

$$w_i(t, x) = \bar{w}_i^\pm, \quad \text{if } X_i^0(Y_i(t, x)) \geq L,$$

namely,

$$w_i(t, x) = \bar{w}_i^+, \quad \forall x \geq X_i^+(t).$$

Similarly,

$$w_i(t, x) = \bar{w}_i^-, \quad \forall x \leq X_i^-(t).$$

From the strict hyperbolicity, we have $X_j^\pm(t) < X_i^\pm(t)$ for all $1 \leq j < i \leq n$. Then

$$w_j(t, x) = \bar{w}_j^+, \quad \forall x \geq X_i^+(t), \quad \forall j \leq i \quad (2.2)$$

and

$$w_j(t, x) = \bar{w}_j^-, \quad \forall x \leq X_i^-(t), \quad \forall j \geq i. \quad (2.3)$$

In particular, we obtain

$$w(t, x) = \bar{w}^+, \quad \forall x \geq X_n^+(t). \quad (2.4)$$

Similarly,

$$w(t, x) = \bar{w}^-, \quad \forall x \leq X_1^-(t). \quad (2.5)$$

Remark 2.1.

(1) For each $i \in K_n$, $x = X_i^\pm(t)$ is the i th characteristic curve issued from the point $(0, \pm L)$, respectively.

Indeed, $X_i^\pm(0) = \pm L$ and it follows from (1.8) that

$$\frac{dX_i^\pm(t)}{dt} = \frac{dX_i(t, Y_i^0(\pm L))}{dt} = \lambda_i(w(t, X_i^\pm)),$$

where w is given by (1.10).

(2) The characteristics $x = X_1^-(t)$ and $x = X_n^+(t)$ are both straight lines.

Indeed, (2.4) and (2.5) imply that

$$\lambda_1(w(t, x)) = \lambda_1(\bar{w}^-) \quad \text{if } x \leq X_1^-(t)$$

and

$$\lambda_n(w(t, x)) = \lambda_n(\bar{w}^+) \quad \text{if } x \geq X_n^+(t).$$

Obviously, $\lambda_1(\bar{w}^-)$ and $\lambda_n(\bar{w}^+)$ are constants, we then obtain

$$X_1^-(t) = \lambda_1(\bar{w}^-)t - L, \quad X_n^+(t) = \lambda_n(\bar{w}^+)t + L.$$

Since the system is linearly degenerate, for $1 \leq i < j \leq n$, two curves $x = X_i^+(t)$ and $x = X_j^-(t)$ intersect only once. Let $t_{i,j} > 0$ be their intersection time. Then $X_i^+(t_{i,j}) = X_j^-(t_{i,j})$ for $1 \leq i < j \leq n$. We denote

$$t_L = \max_{1 \leq i < j \leq n} t_{i,j} \quad (2.6)$$

and

$$D = \{(t, x) \mid x \in \mathbb{R}, t > t_L\}.$$

Noting that the function Y_i^0 is strictly increasing, we deduce from the strict hyperbolicity that

$$X_1^-(t) < X_1^+(t) < X_2^-(t) < X_2^+(t) < \cdots < X_n^-(t) < X_n^+(t), \quad \forall t > t_L. \quad (2.7)$$

Then D can be divided into $2n+1$ sub-domains separated by the characteristics $x = X_i^\pm(t)$ ($i \in K_n$). We denote these sub-domains by D_0 , D_i ($i \in K_n$), D_{n+j} ($j \in K_{n-1}$) and D_{2n} , respectively, where

$$D_0 = \{(t, x) \mid x \leq X_1^-(t), t > t_L\}, \quad D_{2n} = \{(t, x) \mid x > X_n^+(t), t > t_L\},$$

$$D_i = \{(t, x) \mid X_i^-(t) < x \leq X_i^+(t), t > t_L\} \quad (i \in K_n),$$

$$D_{n+i} = \{(t, x) \mid X_i^+(t) < x \leq X_{i+1}^-(t), t > t_L\} \quad (i \in K_{n-1}).$$

From (2.4) and (2.5), we obtain

$$w(t, x) = \bar{w}^-, \quad \forall (t, x) \in D_0$$

and

$$w(t, x) = \bar{w}^+, \quad \forall (t, x) \in D_{2n}.$$

Moreover, (2.2) and (2.3) imply that

$$w(t, x) = \bar{w}^i, \quad \forall (t, x) \in D_{n+i} \quad (i \in K_{n-1}),$$

where

$$\bar{w}^i = (\bar{w}_1^+, \dots, \bar{w}_i^+, \bar{w}_{i+1}^-, \dots, \bar{w}_n^-)^\top, \quad \forall i \in K_n.$$

For $i \in K_n$, from (2.7) we have

$$X_{i-1}^+(t) < X_i^-(t) < x \leq X_{i+1}^-(t) < X_{i+1}^+(t), \quad \forall (t, x) \in D_i,$$

which together with (2.2) and (2.3) gives

$$w_j(t, x) = \bar{w}_j^+, \quad \forall (t, x) \in D_i, \quad \forall j < i$$

and

$$w_j(t, x) = \bar{w}_j^-, \quad \forall (t, x) \in D_i, \quad \forall j > i.$$

Since λ_i is independent of w_i , we obtain

$$\lambda_i(w(t, x)) = \lambda_i(\bar{w}^i), \quad \forall (t, x) \in D_i.$$

Hence, equation (1.1) shows that w_i is a traveling wave in D_i with speed $\lambda_i(\bar{w}^i)$. In view of expression (1.10), we may write

$$w(t, x) = (\bar{w}_1^+, \dots, \bar{w}_{i-1}^+, w_i^0(\phi_i(x - \lambda_i(\bar{w}^i)t)), \bar{w}_{i+1}^-, \dots, \bar{w}_n^-)^\top, \quad \text{if } (t, x) \in D_i, \quad (2.8)$$

where ϕ_i ($i \in K_n$) are Lipschitz functions because X_i^0 and Y_i are Lipschitzian. Remark that X_i^- and X_i^+ , X_i^- and X_j^+ ($i < j$) do not intersect after time t_L . The domains D_i and curves $X_i^\pm$ are illustrated in Figure 1.




Figure 1. Traveling wave solutions after a finite time.

Thus, we get the main result of this section.

Theorem 2.1. *Let the assumptions of Proposition 1.1 hold. Assume furthermore that (2.1) holds. Then there exists a finite time $t_L > 0$ defined by (2.6) and Lipschitz functions ϕ_i ($i \in K_n$) such that*

$$w(t, x) = \begin{cases} (\bar{w}_1^-, \dots, \bar{w}_n^-)^\top, & (t, x) \in D_0, \\ (\bar{w}_1^+, \dots, \bar{w}_{i-1}^+, w_i^0(\phi_i(\eta_i)), \bar{w}_{i+1}^-, \dots, \bar{w}_n^-)^\top, & (t, x) \in D_i, \quad i \in K_n, \\ (\bar{w}_1^+, \dots, \bar{w}_i^+, \bar{w}_{i+1}^-, \dots, \bar{w}_n^-)^\top, & (t, x) \in D_{n+i}, \quad i \in K_{n-1}, \\ (\bar{w}_1^+, \dots, \bar{w}_n^+)^\top, & (t, x) \in D_{2n}, \end{cases}$$

where $\eta_i = x - \lambda_i(\bar{w}^i)t$.

Remark 2.2. *Similar results are still valid under the assumption that system (1.1) has characteristics with constant multiplicity. Indeed, if (1.3) holds, then there are essentially s distinct characteristics μ_p ($p \in K_s$). Similarly to (2.6), we may define a finite time still denoted by $t_L > 0$. In this situation, D is divided into $2s + 1$ sub-domains :*

$$D_p = \{(t, x) \mid X_p^-(t) < x \leq X_p^+(t), \quad t > t_L\}, \quad p \in K_s,$$

$$D_p = \{(t, x) \mid X_p^+(t) < x \leq X_{p+1}^-(t), \quad t > t_L\}, \quad p \in K_{s-1}.$$

On each domain D_{s+p} we have

$$w(t, x) = \bar{w}^{r_p}, \quad (t, x) \in D_{s+p}$$

and

$$w(t, x) = \left(\bar{w}_1^+, \dots, \bar{w}_{r_p-1}^+, w_{r_p-1+1}^0(\phi_p(\eta_p)), \dots, w_{r_p}^0(\phi_p(\eta_p)), \bar{w}_{r_p+1}^-, \dots, \bar{w}_n^- \right)^\top, \quad (t, x) \in D_p,$$

where ϕ_p is a Lipschitz function and $\eta_p = x - \mu_p(\bar{w}^{r_p})t$.

3 Long-time behaviors of entropy solutions

This section is devoted to the study of long-time behaviors of the global entropy solutions in the case that $N_i = N$ for all $i \in K_n$. By means of the explicit expression of the entropy solutions and the special structure of the system, we prove that as t goes to infinity, the entropy solutions converge to traveling wave solutions, which are also of explicit forms in terms of the initial data and the system. More precisely, we assume that there is a constant vector $\bar{w} = (\bar{w}_1, \dots, \bar{w}_n)^\top$ such that $w^0 - \bar{w} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and

$$\lim_{x \rightarrow \pm\infty} w^0(x) = \bar{w}.$$

In particular, if $w^0 - \bar{w} \in W^{1,1}(\mathbb{R})$, the above limits hold. We need also smallness conditions on the initial data in appropriate norms according to the situation $\tilde{\lambda}_i = 0$ or $\tilde{\lambda}_i \neq 0$ (see Lemmas 3.5-3.6). The convergence result holds in the sense of $L^1(\mathbb{R})$ norm (see Theorem 3.1).

Let us denote

$$\bar{\lambda}_i = \lambda_i(\bar{w}_1, \dots, \bar{w}_{i-1}, \bar{w}_{i+1}, \dots, \bar{w}_n), \quad \forall i \in K_n.$$

The first two lemmas are consequences of the structure of the system.

Lemma 3.1. *Under the assumptions of Proposition 1.2, for each $i \in K_n$ we have*

$$\begin{aligned} X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t &= x + \int_{Z^0(x)}^{Z^0(x) + \tilde{\lambda}_i t} \left(\frac{1}{N(\tilde{w}^0(\xi))} - \frac{1}{N(\bar{w})} \right) d\xi \\ &\quad + \int_0^t \left(\frac{M}{N}(\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) - \frac{M}{N}(\bar{w}) \right) d\tau, \end{aligned} \quad (3.1)$$

where $\tilde{w}$ and $\tilde{w}^0$ are defined by (1.15).

Proof. From (1.14) we have

$$\begin{aligned} X(t, Z^0(x) + \tilde{\lambda}_i t) &= \int_0^{Z^0(x) + \tilde{\lambda}_i t} \frac{1}{N(\tilde{w}^0(\xi))} d\xi + \int_0^t \left(\frac{M}{N} \right) (\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) d\tau \\ &= \int_0^{Z^0(x)} \frac{1}{N(\tilde{w}^0(\xi))} d\xi + \int_{Z^0(x)}^{Z^0(x) + \tilde{\lambda}_i t} \frac{1}{N(\tilde{w}^0(\xi))} d\xi \\ &\quad + \int_0^t \left(\frac{M}{N} \right) (\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) d\tau. \end{aligned}$$

A direct computation gives

$$\int_0^{Z^0(x)} \frac{1}{N(\tilde{w}^0(\xi))} d\xi = x.$$

Together with (1.11) at $\bar{w}$, i.e.,

$$\bar{\lambda}_i = \frac{M}{N}(\bar{w}) - \frac{\tilde{\lambda}_i}{N(\bar{w})},$$

we obtain (3.1). □

Lemma 3.2. *It holds*

$$\frac{M}{N} = \frac{\tilde{\lambda}_i \lambda_j - \tilde{\lambda}_j \lambda_i}{\tilde{\lambda}_i - \tilde{\lambda}_j}, \quad \forall i, j \in K_n \text{ such that } \lambda_j \neq \lambda_i. \quad (3.2)$$

In particular,

$$\frac{M}{N} = \lambda_i \quad \text{if } \tilde{\lambda}_i = 0, \quad \forall i \in K_n. \quad (3.3)$$

Proof. From (1.11), we have

$$\tilde{\lambda}_i - N\lambda_i = \tilde{\lambda}_j - N\lambda_j, \quad \forall i, j \in K_n.$$

It follows that

$$\frac{1}{N} = \frac{\lambda_i - \lambda_j}{\tilde{\lambda}_i - \tilde{\lambda}_j}, \quad \forall i, j \in K_n \text{ such that } \lambda_j \neq \lambda_i.$$

Hence, (1.11) yields

$$\frac{M}{N} = \lambda_i - \frac{\tilde{\lambda}_i}{N} = \frac{\tilde{\lambda}_i \lambda_j - \tilde{\lambda}_j \lambda_i}{\tilde{\lambda}_i - \tilde{\lambda}_j}, \quad \forall i, j \in K_n \text{ such that } \lambda_j \neq \lambda_i.$$

This proves the Lemma. $\square$

In the next two lemmas we deal with the limits as $t \rightarrow +\infty$ of the function defined in (3.1). For the sake of simplicity, we only prove the result in the strictly hyperbolic case. In the case of constant multiplicity of eigenvalues, the result can be obtained in an analogous way (see Remark 2.2).

Lemma 3.3. *Let us denote $u = \frac{M}{N}$ and*

$$\psi_{i1}^t(x) = \int_0^t [u(\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) - u(\bar{w})] d\tau, \quad \forall x \in \mathbb{R}, t > 0. \quad (3.4)$$

Let the assumptions of Proposition 1.2 hold and the system be strictly hyperbolic. Let $w^0 - \bar{w} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and $\lim_{x \rightarrow \pm\infty} w^0(x) = \bar{w}$. Then for all $x \in \mathbb{R}$ and $i \in K_n$, we have

$$\lim_{t \rightarrow +\infty} \psi_{i1}^t(x) = \psi_{i1}(x),$$

where

$$\psi_{i1}(x) = \begin{cases} \sum_{\lambda_j > \lambda_i} \frac{1}{\tilde{\lambda}_j - \tilde{\lambda}_i} \int_{-\infty}^{+\infty} (\lambda_i(\bar{w}_1, \dots, \bar{w}_{j-1}, \tilde{w}_j^0(\xi), \bar{w}_{j+1}, \dots, \bar{w}_n) - \bar{\lambda}_i) d\xi \\ + \frac{1}{\tilde{\lambda}_i - \tilde{\lambda}_k} \int_{Z^0(x)}^{+\infty} (\lambda_k(\bar{w}_1, \dots, \bar{w}_{i-1}, \tilde{w}_i^0(\xi), \bar{w}_{i+1}, \dots, \bar{w}_n) - \bar{\lambda}_k) d\xi, & \text{if } \tilde{\lambda}_i > 0, \\ \int_0^{+\infty} (\lambda_i(\tilde{w}(\tau, Z^0(x))) - \lambda_i(\bar{w})) d\tau, & \text{if } \tilde{\lambda}_i = 0, \\ \sum_{\lambda_j < \lambda_i} \frac{1}{\tilde{\lambda}_i - \tilde{\lambda}_j} \int_{-\infty}^{+\infty} (\lambda_i(\bar{w}_1, \dots, \bar{w}_{j-1}, \tilde{w}_j^0(\xi), \bar{w}_{j+1}, \dots, \bar{w}_n) - \bar{\lambda}_i) d\xi \\ + \frac{1}{\tilde{\lambda}_i - \tilde{\lambda}_k} \int_{Z^0(x)}^{-\infty} (\lambda_k(\bar{w}_1, \dots, \bar{w}_{i-1}, \tilde{w}_i^0(\xi), \bar{w}_{i+1}, \dots, \bar{w}_n) - \bar{\lambda}_k) d\xi, & \text{if } \tilde{\lambda}_i < 0, \end{cases} \quad (3.5)$$

with $k \in K_n$ being any index such that $\lambda_k \neq \lambda_i$. Moreover, the above convergence is uniform on any compact set of $\mathbb{R}$.

Proof. Obviously, $\tilde{w}^0 - \bar{w} \in L^1(\mathbb{R})$. Since Z^0 is Lipschitzian and strictly increasing on $\mathbb{R}$, we get $\tilde{w}(\cdot, Z^0(x)) - \bar{w} \in L^1(\mathbb{R})$ for all $x \in \mathbb{R}$. Then the Taylor formula implies that $\lambda_i(\tilde{w}(\cdot, Z^0(x))) - \lambda_i(\bar{w}) \in L^1(\mathbb{R})$. Hence, the result for $\tilde{\lambda}_i = 0$ follows from (3.3).

Now we prove the result for $\tilde{\lambda}_i > 0$ and we omit the proof of the result for $\tilde{\lambda}_i < 0$ since it is similar. First, the Taylor formula gives

$$u(\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) - u(\bar{w}) = \sum_{j=1}^n u_{ij}^t(\tau, x) (\tilde{w}_j(\tau, Z^0(x) + \tilde{\lambda}_i t) - \bar{w}_j),$$

where

$$u_{ij}^t(\tau, x) = \int_0^1 \frac{\partial u}{\partial w_j} (\bar{w} + \alpha(\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t) - \bar{w})) d\alpha.$$

If $\tilde{\lambda}_j = 0$ for some $j \in K_n$, from (3.3) we have $u = \lambda_j$. Since the system is linearly degenerate, λ_j is independent of w_j , so that $\frac{\partial u}{\partial w_j} = 0$. Hence, from (1.15) we may write

$$u(\tilde{w}(\tau, Z^0(x) + \tilde{\lambda}_i t)) - u(\bar{w}) = \sum_{\tilde{\lambda}_j \neq 0} u_{ij}^t(\tau, x) (\tilde{w}_j^0(Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_j \tau) - \bar{w}_j).$$

For each $\tilde{\lambda}_j \neq 0$, we may make a change of variable $\xi = Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_j \tau$ in the integral of the above functions. It follows that

$$\psi_{i1}^t(x) = - \sum_{\tilde{\lambda}_j \neq 0} \frac{1}{\tilde{\lambda}_j} \int_{Z^0(x) + \tilde{\lambda}_i t}^{Z^0(x) + (\tilde{\lambda}_i - \tilde{\lambda}_j)t} u_{ij}^t\left(\frac{Z^0(x) + \tilde{\lambda}_i t - \xi}{\tilde{\lambda}_j}, x\right) (\tilde{w}_j^0(\xi) - \bar{w}_j) d\xi.$$

Note that the function u_{ij}^t is uniformly bounded, i.e., there is a constant $C > 0$ such that

$$|u_{ij}^t(\tau, x)| \leq C, \quad \forall \tau \in [0, t], \quad x \in \mathbb{R}, \quad t > 0.$$

Since $w^0 - \bar{w} \in L^1(\mathbb{R})$, we obtain

$$|u_{ij}^t(\tilde{w}_j^0 - \bar{w}_j)| \leq C |\tilde{w}_j^0 - \bar{w}_j| \in L^1(\mathbb{R}).$$

On the other hand, for all $k \in K_n$, from $\xi = Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_j \tau$ we have

$$Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_k \tau = \frac{\tilde{\lambda}_k}{\tilde{\lambda}_j} \xi + \left(1 - \frac{\tilde{\lambda}_k}{\tilde{\lambda}_j}\right) (Z^0(x) + \tilde{\lambda}_i t).$$

Since Z^0 is a Lipschitz function, for all $x \in \mathbb{R}$ and $\xi \in \mathbb{R}$, we have

$$\lim_{t \rightarrow +\infty} (Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_k \tau) = \infty, \quad \forall k \in K_n \text{ such that } \lambda_k \neq \lambda_j.$$

It follows from (1.15) and $\lim_{x \rightarrow \pm\infty} w^0(x) = \bar{w}$ that

$$\lim_{t \rightarrow +\infty} (\tilde{w}_k^0(Z^0(x) + \tilde{\lambda}_i t - \tilde{\lambda}_k \tau)) = \bar{w}_k, \quad \forall k \in K_n \text{ such that } \lambda_k \neq \lambda_j.$$

Hence,

$$\lim_{t \rightarrow +\infty} u_{ij}^t\left(\frac{Z^0(x) + \tilde{\lambda}_i t - \xi}{\tilde{\lambda}_j}, x\right) = \int_0^1 \frac{\partial u}{\partial w_j} (\bar{w}_1, \dots, \bar{w}_{j-1}, \bar{w}_j + \alpha(\tilde{w}_j^0(\xi) - \bar{w}_j), \bar{w}_{j+1}, \dots, \bar{w}_n) d\alpha.$$

Or

$$\begin{aligned} & \int_0^1 \frac{\partial u}{\partial w_j}(\bar{w}_1, \dots, \bar{w}_{j-1}, \bar{w}_j + \alpha(\tilde{w}_j^0(\xi) - \bar{w}_j), \bar{w}_{j+1}, \dots, \bar{w}_n) d\alpha \cdot (\tilde{w}_j^0(\xi) - \bar{w}_j) \\ &= u(\bar{w}_1, \dots, \bar{w}_{j-1}, \tilde{w}_j^0(\xi), \bar{w}_{j+1}, \dots, \bar{w}_n) - u(\bar{w}). \end{aligned}$$

For $\tilde{\lambda}_j < \tilde{\lambda}_i$, we have $Z^0(x) + (\tilde{\lambda}_i - \tilde{\lambda}_j)t \rightarrow +\infty$. Since $\tilde{\lambda}_i > 0$, we also have $Z^0(x) + \tilde{\lambda}_i t \rightarrow +\infty$. Therefore, for all $x \in \mathbb{R}$, the Lebesgue dominated convergence theorem yields

$$\lim_{t \rightarrow +\infty} \int_{Z^0(x) + \tilde{\lambda}_i t}^{Z^0(x) + (\tilde{\lambda}_i - \tilde{\lambda}_j)t} u_{ij}^t\left(\frac{Z^0(x) + \tilde{\lambda}_i t - \xi}{\tilde{\lambda}_j}, x\right) (\tilde{w}_j^0(\xi) - \bar{w}_j) d\xi = 0.$$

Similarly,

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \int_{Z^0(x) + \tilde{\lambda}_i t}^{Z^0(x) + (\tilde{\lambda}_i - \tilde{\lambda}_j)t} u_{ij}^t\left(\frac{Z^0(x) + \tilde{\lambda}_i t - \xi}{\tilde{\lambda}_j}, x\right) (\tilde{w}_j^0(\xi) - \bar{w}_j) d\xi \\ &= \int_{+\infty}^{-\infty} [u(\bar{w}_1, \dots, \bar{w}_{j-1}, \tilde{w}_j^0(\xi), \bar{w}_{j+1}, \dots, \bar{w}_n) - u(\bar{w})] d\xi, \quad \text{if } \tilde{\lambda}_j > \tilde{\lambda}_i, \end{aligned}$$

and

$$\begin{aligned} & \lim_{t \rightarrow +\infty} \int_{Z^0(x) + \tilde{\lambda}_i t}^{Z^0(x) + (\tilde{\lambda}_i - \tilde{\lambda}_j)t} u_{ij}^t\left(\frac{Z^0(x) + \tilde{\lambda}_i t - \xi}{\tilde{\lambda}_j}, x\right) (\tilde{w}_j^0(\xi) - \bar{w}_j) d\xi \\ &= \int_{+\infty}^{Z^0(x)} [u(\bar{w}_1, \dots, \bar{w}_{i-1}, \tilde{w}_i^0(\xi), \bar{w}_{i+1}, \dots, \bar{w}_n) - u(\bar{w})] d\xi, \quad \text{if } \tilde{\lambda}_j = \tilde{\lambda}_i. \end{aligned}$$

This implies that

$$\lim_{t \rightarrow +\infty} \psi_{i1}^t(x) = \psi_{i1}(x)$$

with

$$\begin{aligned} \psi_{i1}(x) &= \sum_{\tilde{\lambda}_j > \tilde{\lambda}_i} \frac{1}{\tilde{\lambda}_j} \int_{-\infty}^{+\infty} [u(\bar{w}_1, \dots, \bar{w}_{j-1}, \tilde{w}_j^0(\xi), \bar{w}_{j+1}, \dots, \bar{w}_n) - u(\bar{w})] d\xi \\ &\quad + \frac{1}{\tilde{\lambda}_i} \int_{Z^0(x)}^{+\infty} [u(\bar{w}_1, \dots, \bar{w}_{i-1}, \tilde{w}_i^0(\xi), \bar{w}_{i+1}, \dots, \bar{w}_n) - u(\bar{w})] d\xi. \end{aligned}$$

Finally, from (1.11) and $N > 0$, we deduce that $\lambda_i - \lambda_j$ and $\tilde{\lambda}_i - \tilde{\lambda}_j$ have the same sign. Using (3.2) and the fact that λ_j is independent of w_j for all $j \in K_n$, we obtain (3.5) for $\tilde{\lambda}_i > 0$. This proves the pointwise convergence of ψ_{i1}^t on $\mathbb{R}$ for all $i \in K_n$. When x stays in a compact set of $\mathbb{R}$, $Z^0(x)$ is bounded independent of x . This shows that the convergence of ψ_{i1}^t is uniform on any compact set of $\mathbb{R}$. $\square$

Lemma 3.4. *Under the assumptions of Lemma 3.3. For each $i \in K_n$ and all $x \in \mathbb{R}$, we have*

$$\lim_{t \rightarrow +\infty} (X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t) = \psi_i(x), \quad (3.6)$$

where

$$\psi_i(x) = x + \psi_{i1}(x) + \psi_{i2}(x), \quad (3.7)$$

with

$$\psi_{i2}(x) = \begin{cases} \int_{Z^0(x)}^{+\infty} \left(\frac{1}{N(\tilde{w}^0(\xi))} - \frac{1}{N(\bar{w})} \right) d\xi, & \text{if } \tilde{\lambda}_i > 0, \\ 0, & \text{if } \tilde{\lambda}_i = 0, \\ \int_{Z^0(x)}^{-\infty} \left(\frac{1}{N(\tilde{w}^0(\xi))} - \frac{1}{N(\bar{w})} \right) d\xi, & \text{if } \tilde{\lambda}_i < 0. \end{cases} \quad (3.8)$$

Moreover, the convergence (3.6) is uniform on any compact set of $\mathbb{R}$.

Proof. From $w^0 - \bar{w} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$, we have

$$\frac{1}{N(\tilde{w}^0)} - \frac{1}{N(\bar{w})} \in L^1(\mathbb{R}),$$

so that, for all $x \in \mathbb{R}$, when $\tilde{\lambda}_i > 0$,

$$\lim_{t \rightarrow +\infty} \int_{Z^0(x)}^{Z^0(x) + \tilde{\lambda}_i t} \left(\frac{1}{N(\tilde{w}^0)} - \frac{1}{N(\bar{w})} \right) d\xi = \int_{Z^0(x)}^{+\infty} \left(\frac{1}{N(\tilde{w}^0)} - \frac{1}{N(\bar{w})} \right) d\xi.$$

Thus, the result for $\tilde{\lambda}_i > 0$ follows from Lemmas 3.1 and 3.3. The cases $\tilde{\lambda}_i < 0$ and $\tilde{\lambda}_i = 0$ can be treated in a same way. $\square$

The next two results are concerned with regularities and monotonicities of functions ψ_i , $i \in K_n$. To this end, we distinguish again the cases $\tilde{\lambda}_i \neq 0$ and $\tilde{\lambda}_i = 0$. In both cases, we need smallness conditions on the initial data to only assure the invertibility of ψ_i .

Lemma 3.5. *Let the assumptions of Lemma 3.3 hold. For each $i \in K_n$ with $\tilde{\lambda}_i \neq 0$, ψ_i is a Lipschitz function on $\mathbb{R}$. Moreover, there are constants $\delta_1 > 0$ and $C_{\delta_1} > 0$ such that if $\|w^0 - \bar{w}\|_{L^\infty(\mathbb{R})} < \delta_1$, then $\psi'_i(x) \geq C_{\delta_1}$ for almost all $x \in \mathbb{R}$.*

Proof. We only prove the result for $i \in K_n$ with $\tilde{\lambda}_i > 0$. The case for $\tilde{\lambda}_i < 0$ can be treated in a same way.

From (3.5) and (3.7)-(3.8), in the strictly hyperbolic case, we have

$$\begin{aligned} \psi_i(x) &= x + \int_{Z^0(x)}^{+\infty} \left(\frac{1}{N(\tilde{w}^0(\xi))} - \frac{1}{N(\bar{w})} \right) d\xi \\ &\quad + \frac{1}{\tilde{\lambda}_i - \tilde{\lambda}_k} \int_{Z^0(x)}^{+\infty} (\lambda_k(\bar{w}_1, \dots, \bar{w}_{i-1}, \tilde{w}_i^0(\xi), \bar{w}_{i+1}, \dots, \bar{w}_n) - \bar{\lambda}_k) d\xi + \bar{\psi}_i, \end{aligned} \quad (3.9)$$

for any $k \in K_n$ such that $\tilde{\lambda}_k \neq \tilde{\lambda}_i$, where $\bar{\psi}_i$ is a constant. Since Z^0 is a Lipschitz function on $\mathbb{R}$ and the function in the integral of (3.9) is in $L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$, by composition we deduce that ψ_i is a Lipschitz function on $\mathbb{R}$.

Hence, ψ_i is differentiable for almost all $x \in \mathbb{R}$. Let us denote

$$f_i(w) = \frac{1}{N(w)} + \frac{\lambda_k(\bar{w}_1, \dots, \bar{w}_{i-1}, w_i, \bar{w}_{i+1}, \dots, \bar{w}_n)}{\tilde{\lambda}_i - \tilde{\lambda}_k},$$

which is a smooth function. Noting $\tilde{w}^0(Z^0) = w^0$, a straightforward computation yields

$$\psi'_i(x) = 1 - N(w^0(x)) (f_i(w^0(x)) - f_i(\bar{w})).$$

Thus, when $\|w^0 - \bar{w}\|_{L^\infty(\mathbb{R})} < \delta_1$, which is sufficiently small, we obtain $\psi'_i(x) \geq C_{\delta_1} > 0$ for almost all $x \in \mathbb{R}$. This proves the result. $\square$

Lemma 3.6. *Let the assumptions of Lemma 3.3 hold. For each $i \in K_n$ with $\tilde{\lambda}_i = 0$, ψ_i is a uniform continuous function on $\mathbb{R}$. If $w^0 - \bar{w} \in W^{1,1}(\mathbb{R})$, then ψ_i is a Lipschitz function on $\mathbb{R}$. Moreover, there are constants $\delta_2 > 0$ and $C_{\delta_2} > 0$ such that if $\|(w^0)'\|_{L^1(\mathbb{R})} < \delta_2$, then $\psi'_i(x) \geq C_{\delta_2}$ for almost all $x \in \mathbb{R}$.*

Proof. Replacing $w^0 - \bar{w}$ by w^0 , we may suppose $\bar{w} = 0$. Since $\tilde{\lambda}_i = 0$, from (3.5) and (3.7)-(3.8), we have

$$\psi_i(x) = x + \int_0^{+\infty} (\lambda_i(\tilde{w}(\tau, Z^0(x))) - \lambda_i(0)) d\tau.$$

Let $x, y \in \mathbb{R}$. Then

$$\begin{aligned} \psi_i(x) - \psi_i(y) &= (x - y) + \int_0^{+\infty} \left(\lambda_i(\tilde{w}_1^0(Z^0(x) - \tilde{\lambda}_1\tau), \dots, \tilde{w}_n^0(Z^0(x) - \tilde{\lambda}_n\tau)) \right. \\ &\quad \left. - \lambda_i(\tilde{w}_1^0(Z^0(y) - \tilde{\lambda}_1\tau), \dots, \tilde{w}_n^0(Z^0(y) - \tilde{\lambda}_n\tau)) \right) d\tau. \end{aligned}$$

Let us denote

$$\Lambda = \max \left\{ \left| \frac{\partial \lambda_i(w)}{\partial w_j} \right|; w \in L^\infty(\mathbb{R}), \forall i, j \in K_n \right\}.$$

By the Taylor formula, we get

$$|\psi_i(x) - \psi_i(y)| \leq |x - y| + \Lambda \sum_{\tilde{\lambda}_j \neq \tilde{\lambda}_i} \int_0^{+\infty} |\tilde{w}_j^0(Z^0(x) - \tilde{\lambda}_j\tau) - \tilde{w}_j^0(Z^0(y) - \tilde{\lambda}_j\tau)| d\tau.$$

Noting that Z^0 is a Lipschitz function and $w^0 \in L^1(\mathbb{R})$, from the uniform continuity of function $x \mapsto w^0(x + \cdot)$ in $L^1(\mathbb{R})$ (see, for instance, [5]) and by composition, we conclude that ψ_i is uniform continuous on $\mathbb{R}$.

Suppose now $w^0 \in W^{1,1}(\mathbb{R})$. It is easy to check that ψ_i is a Lipschitz function on $\mathbb{R}$ and we have

$$\psi'_i(x) = 1 + \sum_{\tilde{\lambda}_j \neq 0} \int_0^{+\infty} \frac{\partial \lambda_i}{\partial w_j} \cdot \frac{\partial \tilde{w}_j^0(Z^0(x) - \tilde{\lambda}_j\tau)}{\partial x} d\tau.$$

By changes of variables, we get

$$\psi'_i(x) = 1 - \sum_{\tilde{\lambda}_j > 0} \frac{1}{\tilde{\lambda}_j} \int_{Z^0(x)}^{-\infty} \frac{\partial \lambda_i}{\partial w_j} (\tilde{w}_j^0(\xi))' N(w^0(x)) d\xi - \sum_{\tilde{\lambda}_j < 0} \frac{1}{\tilde{\lambda}_j} \int_{Z^0(x)}^{+\infty} \frac{\partial \lambda_i}{\partial w_j} (\tilde{w}_j^0(\xi))' N(w^0(x)) d\xi.$$

Therefore, there is a constant $C_1 > 0$ such that

$$\psi'_i(x) \geq 1 - C \|(w^0)'\|_{L^1(\mathbb{R})}, \quad \forall x \in \mathbb{R}.$$

Thus, when $\|(w^0)'\|_{L^1(\mathbb{R})} < \delta_2$, which is sufficiently small, we get $\psi'_i(x) \geq C_{\delta_2} > 0$ for almost all $x \in \mathbb{R}$. This completes the proof of Lemma 3.6. $\square$

To state and prove the main result of this section, we still need a lemma, which was established by the authors in [23].

Lemma 3.7. *Let $f - \bar{f} \in L^1(\mathbb{R})$ with $\bar{f}$ being a real constant. Let $(e_\nu)_{\nu>0}$ be a sequence of Lipschitz functions defined on $\mathbb{R}$. Suppose that there is a constant $\gamma > 0$ such that $e'_\nu(x) \geq \gamma$, $\forall \nu > 0$, a.e. $x \in \mathbb{R}$ and $e_\nu(x) \rightarrow x$ as $\nu \rightarrow 0$, pointwisely on $\mathbb{R}$ and uniformly on any compact set of $\mathbb{R}$. Then*

$$I_f^\nu = \int_{\mathbb{R}} |f(x) - f(e_\nu(x))| dx \rightarrow 0, \quad \text{as } \nu \rightarrow 0.$$

Theorem 3.1. *Let the assumptions of Lemma 3.3 hold. Furthermore, assume that for each $i \in K_n$, ψ_i is Lipschitzian on $\mathbb{R}$ satisfying $\psi'_i(x) \geq C_0$ for some constant $C_0 > 0$. Let ϕ_i be the inverse function of ψ_i . Then we have*

$$\lim_{t \rightarrow +\infty} \|w_i(t, \cdot) - w_i^0(\phi_i(\cdot - \bar{\lambda}_i t))\|_{L^1(\mathbb{R})} = 0. \quad (3.10)$$

Proof. Let $x = X^0(Z(t, \eta) - \tilde{\lambda}_i t)$. Then $\eta = X(t, Z^0(x) + \tilde{\lambda}_i t)$. From (1.13)-(1.14), we have

$$d\eta = J^0(t, x) dx, \quad \text{with} \quad J^0(t, x) = \frac{N(w^0(x))}{N(\tilde{w}(t, Z^0(x) + \tilde{\lambda}_i t))}.$$

Obviously, there are positive constants denoted by C_1 and C_2 such that $C_1 \leq J^0(t, x) \leq C_2$ for almost all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$. We deduce from the explicit expression of the entropy solutions that

$$\begin{aligned} & \|w_i(t, \cdot) - w_i^0(\phi_i(\cdot - \bar{\lambda}_i t))\|_{L^1(\mathbb{R})} \\ &= \int_{\mathbb{R}} |w_i^0(X^0(Z(t, \eta) - \tilde{\lambda}_i t)) - w_i^0(\phi_i(\eta - \bar{\lambda}_i t))| d\eta \\ &= \int_{\mathbb{R}} |w_i^0(x) - w_i^0(\phi_i(X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t))| J^0(t, x) dx. \end{aligned}$$

Let

$$e_t(x) = \phi_i(X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t).$$

It is easy to see that function $w_i^0 - w_i^0(e_t)$ is uniformly bounded in $L^1(\mathbb{R})$ with respect to t and

$$e'_t(x) = J^0(t, x) \phi'_i(X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t).$$

From the discussion above, there is a positive constant, denoted by γ , such that $e'_t(x) \geq \gamma$ for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$. Since ϕ_i is the inverse function of ψ_i , Lemma 3.4 yields

$$\forall x \in \mathbb{R}, \quad \lim_{t \rightarrow +\infty} e_t(x) = x,$$

pointwisely on $\mathbb{R}$ and uniformly on any compact set of $\mathbb{R}$. Thus, e_t fulfills the conditions of Lemma 3.7 with $\Omega_1 = \mathbb{R}$ and $\nu = \frac{1}{t}$. We conclude that

$$\lim_{t \rightarrow +\infty} \int_{\mathbb{R}} |w_i^0(x) - w_i^0(\phi_i(X(t, Z^0(x) + \tilde{\lambda}_i t) - \bar{\lambda}_i t))| J^0(t, x) dx = 0,$$

which completes the proof of Theorem 3.1. $\square$

Remark 3.1. *In Theorem 3.1, the assumption $\psi'_i(x) \geq C_0$ for all $x \in \mathbb{R}$ is essential. It implies that ψ_i is a strictly increasing function on $\mathbb{R}$. Sufficient conditions to ensure this assumption are given in Lemmas 3.5 and 3.6, from which we see that the case $\tilde{\lambda}_i = 0$ requires more regularity on the initial data. In particular, if $\tilde{\lambda}_i \neq 0$ for some $i \in K_n$, we only require that $w^0 - \bar{w} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ and the L^∞ norm of $w^0 - \bar{w}$ is small to yield (3.10).*

4 L^1 stability of entropy solutions

This section is concerned with the L^1 stability of entropy solutions to the Cauchy problem (1.1)-(1.2) in the case $N_i = N$ for all $i \in K_n$. To this end, suppose $w^{1,0}, w^{2,0} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$

and let w^1 and w^2 be the entropy solutions with initial data $w^{1,0}$ and $w^{2,0}$, respectively. For simplicity, we define

$$R_0 = \|w^{2,0} - w^{1,0}\|_{L^1(\mathbb{R})} \quad \text{and} \quad R_t = \|w^2(t, \cdot) - w^1(t, \cdot)\|_{L^1(\mathbb{R})}.$$

Here the L^1 stability of the entropy solutions to the Cauchy problem (1.1)-(1.2) means that for all $\varepsilon > 0$, there exists $\delta > 0$ depending only on ε such that

$$R_0 \leq \delta \implies R_t \leq \varepsilon, \quad \forall t > 0,$$

or equivalently,

$$\lim_{R_0 \rightarrow 0} R_t = 0, \quad \forall t > 0. \quad (4.1)$$

Together with the uniqueness of entropy solutions, stability condition (4.1) implies that application $R_0 \mapsto R_t$ is continuous at $R_0 = 0$ for all $t \geq 0$. In addition, the strong L^1 stability means that the following stability inequality holds for some constant $C > 0$ independent of t :

$$R_t \leq C R_0, \quad \forall t > 0. \quad (4.2)$$

The study of the strong L^1 stability needs usually to work with small BV initial data, see [4] for a general system of conservation laws and [7] for system (1.1).

For each $j = 1, 2$, since $w^{j,0} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$, from Proposition 1.2, the Cauchy problem (1.1)-(1.2) with the initial data $w^{j,0}$ admits a unique entropy solution, given by

$$w^j(t, x) = \left(w_1^{j,0}(X^{j,0}(Z^j(t, x) - \tilde{\lambda}_1 t)), \dots, w_n^{j,0}(X^{j,0}(Z^j(t, x) - \tilde{\lambda}_n t)) \right)^\top,$$

where $X^{j,0}$ is the inverse function of $Z^{j,0}$ and $Z^j(t, \cdot)$ is the inverse function of $X^j(t, \cdot)$, in which

$$Z^{j,0}(x) = \int_0^x N(w^{j,0}(\xi)) \, d\xi$$

and

$$dX^j(t, z) = \frac{1}{N(\tilde{w}^j(t, z))} \, dz + \frac{M(\tilde{w}^j(t, z))}{N(\tilde{w}^j(t, z))} \, dt,$$

with

$$\tilde{w}^j(t, z) = \left(w_1^{j,0}(X^{j,0}(z - \tilde{\lambda}_1 t)), \dots, w_n^{j,0}(X^{j,0}(z - \tilde{\lambda}_n t)) \right)^\top. \quad (4.3)$$

The main result of this section is as follows.

Theorem 4.1. *Under the assumptions of Proposition 1.2, the unique global entropy solution to the Cauchy problem (1.1)-(1.2) is L^1 stable. If furthermore, $w^{j,0} \in W^{1,1}(\mathbb{R})$ for $j = 1, 2$, then, for each $i \in K_n$, we have*

$$\|w_i^2(t, \cdot) - w_i^1(t, \cdot)\|_{L^1(\mathbb{R})} \leq C R_0, \quad \forall t \geq 0. \quad (4.4)$$

The proof of Theorem 4.1 is based on a series of Lemmas below.

Lemma 4.1. *Let the assumptions of Proposition 1.2 hold. There exists a positive constant C , such that for all $x, z \in \mathbb{R}$, we have*

$$|Z^{2,0}(x) - Z^{1,0}(x)| \leq C R_0, \quad |X^{2,0}(z) - X^{1,0}(z)| \leq C R_0$$

and

$$|X^{1,0}(Z^{2,0}(x)) - x| \leq C R_0. \quad (4.5)$$

Proof. The proof is omitted here since it is similar to that of Lemma 4.1 in [23]. $\square$

The next lemma was proved by the authors in [23].

Lemma 4.2. *Let $f \in L^1_{loc}(\mathbb{R})$, $f' \in L^1(\mathbb{R})$ and h_t be a Lipschitz function on $\mathbb{R}$ such that*

$$\text{a.e. } (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \quad h'_t(x) \geq \gamma_1, \quad |h_t(x) - x| \leq \gamma_2(t),$$

for a constant $\gamma_1 > 0$ and a function $\gamma_2(t) \geq 0$. Then $f - f(h_t) \in L^1(\mathbb{R})$ and we have

$$\int_{\mathbb{R}} |f(x) - f(h_t(x))| dx \leq \frac{\gamma_2(t)}{\gamma'_1} \|f'\|_{L^1(\mathbb{R})},$$

where $\gamma'_1 = \min\{1, \gamma_1\}$.

Lemma 4.3. *Under the assumptions of Proposition 1.2, we have*

$$|X^2 - X^1| \longrightarrow 0 \quad \text{and} \quad |Z^2 - Z^1| \longrightarrow 0, \quad \text{as} \quad R_0 \rightarrow 0, \quad (4.6)$$

uniformly on $\mathbb{R}^+ \times \mathbb{R}$. If furthermore, $w^{j,0} \in W^{1,1}(\mathbb{R})$ for $j = 1, 2$, then, for all $t \geq 0$ and all $x, z \in \mathbb{R}$, we have

$$|X^2(t, z) - X^1(t, z)| \leq C R_0, \quad |Z^2(t, x) - Z^1(t, x)| \leq C R_0. \quad (4.7)$$

Proof. For $x \in \mathbb{R}$, let $z_2 = Z^2(t, x)$. Then $x = X^2(t, z_2)$. Since $Z^1(t, \cdot)$ is Lipschitzian on $\mathbb{R}$, we have

$$\begin{aligned} |Z^2(t, x) - Z^1(t, x)| &= |Z^2(t, X^2(t, z_2)) - Z^1(t, X^2(t, z_2))| \\ &= |Z^1(t, X^1(t, z_2)) - Z^1(t, X^2(t, z_2))| \\ &\leq C |X^2(t, z_2) - X^1(t, z_2)|. \end{aligned}$$

Hence, the second estimate of (4.6) immediately follows from the first one. It remains to prove the first estimate of (4.6).

From (1.17), for any $i \in K_n$ we have

$$X^j(t, z) = \int_0^z \frac{d\xi}{N(w^{j,0}(X^{j,0}(\xi)))} + \int_0^t \lambda_i(\tilde{w}^j(\tau, z)) d\tau - \int_0^t \frac{\tilde{\lambda}_i d\tau}{N(\tilde{w}^j(\tau, z))},$$

where $\tilde{w}^j(t, z)$ ($j = 1, 2$) is defined by (4.3). Therefore

$$\begin{aligned} |X^2(t, z) - X^1(t, z)| &\leq \int_{\mathbb{R}} \left| \frac{1}{N(w^{2,0}(X^{2,0}(\xi)))} - \frac{1}{N(w^{1,0}(X^{1,0}(\xi)))} \right| d\xi \\ &\quad + \int_{\mathbb{R}^+} |\lambda_i(\tilde{w}^2(\tau, z)) - \lambda_i(\tilde{w}^1(\tau, z))| d\tau \\ &\quad + \tilde{\lambda}_i \int_{\mathbb{R}^+} \left| \frac{1}{N(\tilde{w}^2(\tau, z))} - \frac{1}{N(\tilde{w}^1(\tau, z))} \right| d\tau \\ &\stackrel{\text{def}}{=} \mathcal{F} + \mathcal{G}_i + \mathcal{H}_i. \end{aligned}$$

Since $N > 0$, we have

$$\mathcal{F} = \int_{\mathbb{R}} \frac{|N(w^{2,0}(X^{2,0}(\xi))) - N(w^{1,0}(X^{1,0}(\xi)))|}{N(w^{1,0}(X^{1,0}(\xi))) \cdot N(w^{2,0}(X^{2,0}(\xi)))} d\xi$$

$$\begin{aligned}
&\leq C \int_{\mathbb{R}} |w^{2,0}(X^{2,0}(\xi)) - w^{1,0}(X^{1,0}(\xi))| \, d\xi \\
&\leq C \int_{\mathbb{R}} |w^{2,0}(X^{2,0}(\xi)) - w^{1,0}(X^{2,0}(\xi))| \, d\xi \\
&\quad + C \int_{\mathbb{R}} |w^{1,0}(x) - w^{1,0}(X^{1,0}(Z^{2,0}(x)))| \, dx.
\end{aligned} \tag{4.8}$$

It follows from (4.5) that

$$X^{1,0}(Z^{2,0}(x)) \longrightarrow x, \quad \text{as } R_0 \rightarrow 0,$$

uniformly on $\mathbb{R}$. Moreover,

$$\forall x \in \mathbb{R}, \quad \left(X^{1,0}(Z^{2,0}(x)) \right)' = \frac{N(w^{2,0}(x))}{N(w^{1,0}(X^{1,0}(Z^{2,0}(x))))} \geq \gamma, \tag{4.9}$$

for some constant $\gamma > 0$. We conclude from Lemma 3.7 with $\Omega_1 = \mathbb{R}$ that

$$\int_{\mathbb{R}} |w^{1,0}(x) - w^{1,0}(X^{1,0}(Z^{2,0}(x)))| \, dx \longrightarrow 0, \quad \text{as } R_0 \rightarrow 0,$$

which together with (4.8) implies that $\mathcal{F} \longrightarrow 0$ as $R_0 \rightarrow 0$.

Next, we consider $\mathcal{G}_i$ and $\mathcal{H}_i$. If $\tilde{\lambda}_k \neq 0$ for all $k \in K_n$, then for any fixed $i \in K_n$, $\mu_i = \lambda_i$ or $\mu_i = \tilde{\lambda}_i/N$, we have

$$\begin{aligned}
&\int_{\mathbb{R}^+} |\mu_i(\tilde{w}^2(\tau, z)) - \mu_i(\tilde{w}^1(\tau, z))| \, d\tau \\
&\leq C \sum_{k=1}^n \int_{\mathbb{R}^+} \left| w_k^{2,0}(X^{2,0}(z - \tilde{\lambda}_k \tau)) - w_k^{1,0}(X^{1,0}(z - \tilde{\lambda}_k \tau)) \right| \, d\tau,
\end{aligned}$$

which can be treated as for $\mathcal{F}$ due to changes of variables $\tau \mapsto z - \tilde{\lambda}_k \tau$. This shows $\mathcal{G}_i \longrightarrow 0$ and $\mathcal{H}_i \longrightarrow 0$ as $R_0 \rightarrow 0$. If $\tilde{\lambda}_i = 0$ for some $i \in K_n$, then $\mathcal{H}_i = 0$. Since λ_i is linearly degenerate, λ_i is independent of all w_k such that $\lambda_k = \lambda_i$. Therefore,

$$\begin{aligned}
\mathcal{G}_i &\stackrel{\text{def}}{=} \int_{\mathbb{R}^+} |\lambda_i(\tilde{w}^2(\tau, z)) - \lambda_i(\tilde{w}^1(\tau, z))| \, d\tau \\
&\leq \sum_{\tilde{\lambda}_k \neq 0}^n \int_{\mathbb{R}^+} \left| \frac{\partial \lambda_i}{\partial w_k} \right| \cdot \left| w_k^{2,0}(X^{2,0}(z - \tilde{\lambda}_k \tau)) - w_k^{1,0}(X^{1,0}(z - \tilde{\lambda}_k \tau)) \right| \, d\tau \\
&\leq C \sum_{\tilde{\lambda}_k \neq 0}^n \int_{\mathbb{R}^+} \left| w_k^{2,0}(X^{2,0}(z - \tilde{\lambda}_k \tau)) - w_k^{1,0}(X^{1,0}(z - \tilde{\lambda}_k \tau)) \right| \, d\tau.
\end{aligned}$$

Hence, it is still possible to make the change of variables $\tau \mapsto z - \tilde{\lambda}_k \tau$ to obtain $\mathcal{G}_i \longrightarrow 0$ as $R_0 \rightarrow 0$. This proves (4.6) for X^1 and X^2 .

Finally, similarly to the proof of Lemma 4.2 in [23], noting that (4.5) and (4.9), we can apply Lemma 4.2 to get the desired (4.7). Thus, the proof of Lemma 4.3 is completed. $\square$

Lemma 4.4. *Under the assumptions of Proposition 1.2, we define*

$$\forall (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \quad g_t(x) = X^{1,0}(Z^1(t, X^2(t, Z^{2,0}(x) + \tilde{\lambda}_i t)) - \tilde{\lambda}_i t).$$

Then there is a constant $\gamma > 0$ such that

$$\text{a.e. } (t, x) \in \mathbb{R}^+ \times \mathbb{R}, \quad \frac{dg_t(x)}{dx} \geq \gamma \quad \text{and} \quad g_t(x) \longrightarrow x, \quad \text{as } R_0 \rightarrow 0, \tag{4.10}$$

where the convergence is uniform on $\mathbb{R}^+ \times \mathbb{R}$. If furthermore, $w^{j,0} \in W^{1,1}(\mathbb{R})$ for $j = 1, 2$, then

$$|g_t(x) - x| \leq CR_0, \quad \forall t \geq 0, \quad \forall x \in \mathbb{R}. \quad (4.11)$$

Proof. The proof is omitted here since it is similar to that of Lemma 4.3 in [23]. $\square$

Proof of Theorem 4.1. It suffices to establish the L^1 stability for each w_i ($i \in K_n$). From the explicit expression of the entropy solution, we have

$$\begin{aligned} & \|w_i^2(t, \cdot) - w_i^1(t, \cdot)\|_{L^1(\mathbb{R})} \\ &= \int_{\mathbb{R}} \left| w_i^{2,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{1,0}(Z^1(t, x) - \tilde{\lambda}_i t)) \right| dx \\ &\leq \int_{\mathbb{R}} \left| w_i^{2,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) \right| dx \\ &\quad + \int_{\mathbb{R}} \left| w_i^{1,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{1,0}(Z^1(t, x) - \tilde{\lambda}_i t)) \right| dx. \end{aligned} \quad (4.12)$$

Using the change of variable

$$\eta_i = X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t),$$

we get

$$x = X^2(t, Z^{2,0}(\eta_i) + \tilde{\lambda}_i t) \quad \text{and} \quad dx = \bar{J}^{2,0}(t, \eta_i) d\eta_i,$$

where

$$\bar{J}^{2,0}(t, \eta_i) = \frac{N(w^{2,0}(\eta_i))}{N(w^{2,0}(X^{2,0}(Z^{2,0}(\eta_i) + \tilde{\lambda}_i t)))} \geq \gamma > 0, \quad (4.13)$$

with γ being a constant. It is easy to see that

$$\begin{aligned} & \int_{\mathbb{R}} \left| w_i^{2,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) \right| dx \\ &= \int_{\mathbb{R}} |w_i^{2,0}(\eta_i) - w_i^{1,0}(\eta_i)| \bar{J}^{2,0}(t, \eta_i) d\eta_i \\ &\leq C \|w_i^{2,0} - w_i^{1,0}\|_{L^1(\mathbb{R})}. \end{aligned} \quad (4.14)$$

For the second term of the right hand side of (4.12), we have

$$\begin{aligned} & \int_{\mathbb{R}} \left| w_i^{1,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{1,0}(Z^1(t, x) - \tilde{\lambda}_i t)) \right| dx \\ &= \int_{\mathbb{R}} |w_i^{1,0}(\eta_i) - w_i^{1,0}(g_t(\eta_i))| \bar{J}^{2,0}(t, \eta_i) d\eta_i. \end{aligned} \quad (4.15)$$

Together with $w_i^{1,0} \in L^1(\mathbb{R})$ and (4.10), we may apply Lemma 3.7 to get

$$\int_{\mathbb{R}} \left| w_i^{1,0}(X^{2,0}(Z^2(t, x) - \tilde{\lambda}_i t)) - w_i^{1,0}(X^{1,0}(Z^1(t, x) - \tilde{\lambda}_i t)) \right| dx \longrightarrow 0, \quad \text{as } R_0 \rightarrow 0.$$

This shows that

$$\|w_i^2(t, \cdot) - w_i^1(t, \cdot)\|_{L^1(\mathbb{R})} \longrightarrow 0, \quad \text{as } R_0 \rightarrow 0.$$

If $w^{j,0} \in W^{1,1}(\mathbb{R})$ for $j = 1, 2$, with the aid of (4.11), we may apply Lemma 4.2 to (4.15). Thus, together with (4.12)-(4.15), it gives (4.4). $\square$

5 Examples

We present two examples for the results obtained in sections 2-4. They are the Born-Infeld system (of 4 equations) and the augmented Born-Infeld system (of 8 equations) introduced by Brenier [3], see also [26]. We mention that both the Born-Infeld system and the augmented Born-Infeld system are rich and linearly degenerate with constant multiplicity eigenvalues. Moreover, they have a common structure : $N_i = N$ for all $i \in K_n$ (see [16]). Another example is the generalized extremal surface equations, which have been analyzed in detail in [23].

5.1 The Born-Infeld system

The one-dimensional Born-Infeld system reads (see [3]) :

$$\left\{ \begin{array}{l} \partial_t D_2 + \partial_x \left(\frac{B_3 + D_2 P_1 - D_1 P_2}{h} \right) = 0, \\ \partial_t D_3 + \partial_x \left(\frac{-B_2 + D_3 P_1 - D_1 P_3}{h} \right) = 0, \\ \partial_t B_2 + \partial_x \left(\frac{-D_3 + B_2 P_1 - B_1 P_2}{h} \right) = 0, \\ \partial_t B_3 + \partial_x \left(\frac{D_2 + B_3 P_1 - B_1 P_3}{h} \right) = 0, \\ P(u) = D \times B, \quad h(u) = \sqrt{1 + |B|^2 + |D|^2 + |D \times B|^2}. \end{array} \right. \quad (5.1)$$

Here $u = (D_2, D_3, B_2, B_3)^\top$ are the unknown variables, B_1, D_1 are real constants, and

$$B = (B_1, B_2, B_3)^\top, \quad D = (D_1, D_2, D_3)^\top, \quad P = (P_1, P_2, P_3)^\top.$$

From [22], system (5.1) can be written in the form (1.1). It is not strictly hyperbolic with two eigenvalues, denoted by λ and μ , of both constant multiplicity 2. The variables λ and μ satisfy

$$\left\{ \begin{array}{l} \partial_t \mu + \lambda \partial_x \mu = 0, \\ \partial_t \lambda + \mu \partial_x \lambda = 0. \end{array} \right. \quad (5.2)$$

Moreover, system (5.1) admits an additional conservation law (see [3]) :

$$\partial_t h + \partial_x P_1 = 0.$$

From [16], (5.1) is a linearly degenerate rich system with

$$M = P_1, \quad N_i = N \stackrel{def}{=} h = \frac{2a}{\mu - \lambda}, \quad \mu > \lambda.$$

Here N_i is independent of i for all $i \in K_n$. The eigenvalues in Lagrangian coordinates are

$$\tilde{\lambda}_1 = \tilde{\lambda}_2 = -a, \quad \tilde{\lambda}_3 = \tilde{\lambda}_4 = a, \quad (5.3)$$

where $a = \sqrt{1 + B_1^2 + D_1^2}$.

Theorems 3.1 and 4.1 can be applied to the Born-Infeld system. More precisely, given an initial condition for (5.1) :

$$t = 0 : \quad u = u^0(x), \quad x \in \mathbb{R}, \quad (5.4)$$

with $u^0 - \bar{u} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ for some constant $\bar{u} \in \mathbb{R}^4$. We infer from Theorem 4.1 that the unique entropy solution of (5.1) and (5.4) is L^1 stable. Moreover, if $u^0 - \bar{u} \in W^{1,1}(\mathbb{R})$, we get the strong stability inequality like (4.2). In addition, when the initial data satisfy the assumptions of Theorem 3.1, the explicit asymptotic limit can be obtained. In this particular case, we can present better characterization of the explicit asymptotic limit.

For this purpose, we denote $\lambda^0 = \lambda(u^0)$ and $\mu^0 = \mu(u^0)$ and suppose

$$\inf_{x \in \mathbb{R}} \mu^0(x) > \sup_{x \in \mathbb{R}} \lambda^0(x). \quad (5.5)$$

Define

$$Z^0(x) \stackrel{\text{def}}{=} \int_0^x h(u^0(\xi)) d\xi = \int_0^x \frac{2a}{\mu^0(\xi) - \lambda^0(\xi)} d\xi$$

and X^0 being the inverse function of Z^0 . It follows from (1.17) and $X(0, z) = X^0(z)$ that

$$X(t, z) = \frac{1}{2a} \int_0^{z+at} \mu^0(X^0(\xi)) d\xi - \frac{1}{2a} \int_0^{z-at} \lambda^0(X^0(\xi)) d\xi. \quad (5.6)$$

This formula was first obtained in [21]. Suppose now

$$\lim_{x \rightarrow +\infty} \mu^0(x) = \bar{\mu}_+^0, \quad \lim_{x \rightarrow -\infty} \lambda^0(x) = \bar{\lambda}_-^0, \quad (5.7)$$

where $\bar{\mu}_+^0 = \mu(\bar{u})$ and $\bar{\lambda}_-^0 = \lambda(\bar{u})$.

Remark 5.1. *Assumption (5.7) requires the limit only at one direction for each variable. It is weaker than $\lim_{x \rightarrow \pm\infty} w^0(x) = \bar{w}$, which is made in Theorem 3.1 and Lemmas 3.3-3.6.*

Let H and I be the primitives of functions $y \mapsto \lambda^0(X^0(y)) - \bar{\lambda}_-^0$ and $y \mapsto \mu^0(X^0(y)) - \bar{\mu}_+^0$, defined by

$$H(y) = \int_1^y (\lambda^0(X^0(\xi)) - \bar{\lambda}_-^0) d\xi, \quad I(y) = \int_1^y (\mu^0(X^0(\xi)) - \bar{\mu}_+^0) d\xi, \quad \forall y \in \mathbb{R}. \quad (5.8)$$

Since $H' \in L^1(\mathbb{R})$ and $I' \in L^1(\mathbb{R})$, there are two constants $\bar{H}^-$ and $\bar{I}^+$ such that

$$\lim_{y \rightarrow -\infty} H(y) = \bar{H}^-, \quad \lim_{y \rightarrow +\infty} I(y) = \bar{I}^+. \quad (5.9)$$

Define

$$\psi(x) = x + \frac{1}{2a} \left(H(Z^0(x)) - \bar{H}^- \right), \quad \varphi(x) = x - \frac{1}{2a} \left(I(Z^0(x)) - \bar{I}^+ \right), \quad \forall x \in \mathbb{R}. \quad (5.10)$$

A straightforward computation shows that

$$\psi'(x) = \frac{\mu^0(x) - \bar{\lambda}_-^0}{\mu^0(x) - \lambda^0(x)} > 0, \quad \varphi'(x) = \frac{\bar{\mu}_+^0 - \lambda^0(x)}{\mu^0(x) - \lambda^0(x)} > 0.$$

Therefore, ψ and φ are all Lipschitz and strictly increasing functions on $\mathbb{R}$ without any smallness condition on u^0 . Let us denote by ϕ and χ be the inverse functions of ψ and φ , respectively, which are both Lipschitzian and strictly increasing on $\mathbb{R}$.

It follows from (5.6) and (5.9)-(5.10) that, for all $x \in \mathbb{R}$,

$$X(t, Z^0(x) - at) - \bar{\lambda}_-^0 t$$

$$\begin{aligned}
&= \frac{1}{2a} \int_0^{Z^0(x)} \mu^0(X^0(\xi)) d\xi - \frac{1}{2a} \int_0^{Z^0(x)-2at} \lambda^0(X^0(\xi)) d\xi - \bar{\lambda}_-^0 t \\
&= x + \frac{1}{2a} \left(H(Z^0(x)) - H(Z^0(x) - 2at) \right) \\
&= \psi(x) - \frac{1}{2a} \left(H(Z^0(x) - 2at) - \bar{H}^- \right) \\
&\longrightarrow \psi(x), \quad \text{as } t \rightarrow +\infty,
\end{aligned}$$

which implies that

$$\phi(X(t, Z^0(x) - at) - \bar{\lambda}_-^0 t) \longrightarrow x, \quad \text{as } t \rightarrow +\infty, \quad (5.11)$$

pointwisely on $\mathbb{R}$ and uniformly on any compact set of $\mathbb{R}$. Similarly,

$$\chi(X(t, Z^0(x) + at) - \bar{\mu}_+^0 t) \longrightarrow x, \quad \text{as } t \rightarrow +\infty,$$

pointwisely on $\mathbb{R}$ and uniformly on any compact set of $\mathbb{R}$. Let w_1 and w_2 be the Riemann invariants associated to the eigenvalue λ , w_3 and w_4 be those associated to μ (see [22] for the detailed expressions). Then the asymptotic limit of w_i ($1 \leq i \leq 4$) is

$$\begin{cases} \|w_j(t, \cdot) - w_j^0(\phi(\cdot - \bar{\lambda}_-^0 t))\|_{L^1(\mathbb{R})} \longrightarrow 0, & \text{as } t \rightarrow +\infty, \\ \|w_{j+2}(t, \cdot) - w_{j+2}^0(\chi(\cdot - \bar{\mu}_+^0 t))\|_{L^1(\mathbb{R})} \longrightarrow 0, & \text{as } t \rightarrow +\infty, \end{cases} \quad (5.12)$$

for $j = 1, 2$.

From (5.3), no eigenvalue of the Born-Infeld system in Lagrangian coordinates is zero. Applying Theorem 3.1 to (5.1) together with the discussion above, we obtain the following result.

Corollary 5.1. *Let $u^0 - \bar{u} \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. Assume (5.5) and (5.7) hold. Then the entropy solution of (5.1) and (5.4) satisfies (5.12).*

5.2 The augmented Born-Infeld system

The augmented Born-Infeld system reads (see [3]) :

$$\left\{ \begin{aligned} &\partial_t h + \partial_x P_1 = 0, \\ &\partial_t P_1 + \partial_x \left(\frac{P_1^2 - a^2}{h} \right) = 0, \\ &\partial_t D_2 + \partial_x \left(\frac{B_3 + D_2 P_1 - D_1 P_2}{h} \right) = 0, \\ &\partial_t D_3 + \partial_x \left(\frac{-B_2 + D_3 P_1 - D_1 P_3}{h} \right) = 0, \\ &\partial_t B_2 + \partial_x \left(\frac{-D_3 + B_2 P_1 - B_1 P_2}{h} \right) = 0, \\ &\partial_t B_3 + \partial_x \left(\frac{D_2 + B_3 P_1 - B_1 P_3}{h} \right) = 0, \\ &\partial_t P_2 + \partial_x \left(\frac{P_1 P_2 - D_1 D_2 - B_1 B_2}{h} \right) = 0, \\ &\partial_t P_3 + \partial_x \left(\frac{P_1 P_3 - D_1 D_3 - B_1 B_3}{h} \right) = 0, \end{aligned} \right. \quad (5.13)$$

where $h, P_1, D_2, D_3, B_2, B_3, P_2$ and P_3 are the unknown variables. From [22], the eigenvalues (5.13) are :

$$\lambda \stackrel{\text{def}}{=} \lambda_1 = \lambda_2 = \lambda_3 = \frac{P_1 - a}{h}, \quad \nu \stackrel{\text{def}}{=} \lambda_4 = \lambda_5 = \frac{P_1}{h}, \quad \mu \stackrel{\text{def}}{=} \lambda_6 = \lambda_7 = \lambda_8 = \frac{P_1 + a}{h}.$$

It is easy to see that λ and μ are also the Riemann invariants satisfying (5.2). Moreover, the eigenvalues of system (5.13) in Lagrangian coordinates are :

$$\tilde{\lambda}_1 = \tilde{\lambda}_2 = \tilde{\lambda}_3 = -a, \quad \tilde{\lambda}_4 = \tilde{\lambda}_5 = 0, \quad \tilde{\lambda}_6 = \tilde{\lambda}_7 = \tilde{\lambda}_8 = a.$$

Since the augmented Born-Infeld system and the Born-Infeld system have a common structure : $N_i = h$ ($1 \leq i \leq 8$) and the same eigenvalues λ and μ , it suffices to consider the case for $\tilde{\lambda}_4 = \tilde{\lambda}_5 = 0$. According to the discussion above, we only need to check that the expression like (5.11) holds for ν .

Indeed, with the notations of the previous subsection, we have

$$\begin{aligned} & X(t, Z^0(x)) - \frac{\bar{\lambda}_-^0 + \bar{\mu}_+^0}{2} t \\ &= \frac{1}{2a} \int_0^{Z^0(x)+at} \mu^0(X^0(\xi)) d\xi - \frac{1}{2a} \int_0^{Z^0(x)-at} \lambda^0(X^0(\xi)) d\xi - \frac{\bar{\lambda}_-^0 + \bar{\mu}_+^0}{2} t \\ &= x + \frac{1}{2a} \left(I(Z^0(x) + at) - I(Z^0(x)) + \frac{1}{2a} \left(H(Z^0(x)) - H(Z^0(x) - at) \right) \right) \\ &\longrightarrow x + \frac{1}{2a} \left((H(Z^0(x)) - \bar{H}^-) - (I(Z^0(x)) - \bar{I}^+) \right) \stackrel{\text{def}}{=} \omega(x), \quad \text{as } t \rightarrow +\infty, \end{aligned}$$

where we have used the assumption (5.9). A straightforward calculation shows that

$$\omega'(x) = \frac{\bar{\mu}_+^0 - \bar{\lambda}_-^0}{\mu^0(x) - \lambda^0(x)} > 0.$$

Then we can define its inverse function denoted by θ , which is also a Lipschitz and strictly increasing function on $\mathbb{R}$. Moreover,

$$\theta \left(X(t, Z^0(x)) - \frac{\bar{\lambda}_-^0 + \bar{\mu}_+^0}{2} t \right) \longrightarrow x, \quad \text{as } t \rightarrow +\infty,$$

pointwisely on $\mathbb{R}$ and uniformly on any compact set of $\mathbb{R}$.

Thus, for the eigenvalue ν which does not appear in the Born-Infeld system, there still exists a Lipschitz function θ such that the expression like (5.11) holds. This shows that for initial data in $L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$ the entropy solution of the augmented Born-Infeld system has analogous asymptotic limits to (5.12), without any smallness condition on the initial data. The L^1 stability of solutions can be obtained in a same way as for the Born-Infeld system.

Acknowledgements

This work was in part carried out when Yong-Fu YANG had a postdoctor position of the academic year 2009-2010 at Université Blaise Pascal. He would like to thank Laboratoire de Mathématiques (CNRS-UMR 6620) for hospitality. His research was also supported in part by National Natural Science Foundation of China under Grant No. 10926162, Fundamental Research Funds for the Central Universities No. 2009B01314 and Natural Science Foundation of Hohai University under Grant No. 2009428011.

Références

- [1] F.Bouchut and B.Perthame, Kruzkov's estimates for scalar conservation laws revisited, *Trans. Amer. Math. Soc.* 350 (1998), no. 7, 2847-2870.
- [2] Y. Brenier, Some geometric PDEs related to hydrodynamics and electrodynamics, *Proceedings of the International Congress of Mathematicians*, Vol. 3, (Higher Education Press, Beijing, 2002), 761-772.
- [3] Y. Brenier, Hydrodynamic structure of the augmented Born-Infeld equations, *Arch. Rat. Mech. Anal.* 172 (2004), 65-91.
- [4] A. Bressan, Hyperbolic Systems of Conservation Laws : The One Dimensional Cauchy Problem, Oxford Lecture Series in Math. and its Applications, Vol. 20, 2000.
- [5] H.Brézis, Analyse Fonctionnelle, Masson, Paris, 1993.
- [6] D. Chae and H. Huh, Global existence for small initial data in the Born-Infeld equations, *J. Math. Phys.* 44 (2004), 6132-6139.
- [7] G.Q. Chen, The method of quasidecoupling for discontinuous solutions to conservation laws, *Arch. Rational Mech. Anal.* 172 (1992), 131-185.
- [8] J. Glimm, Solutions in the large for nonlinear hyperbolic systems of equations, *Comm. Pure Appl. Math.*, 18 (1965), 695-715.
- [9] J. Glimm and P.D. Lax, Decay of solutions of system of nonlinear hyperbolic conservation laws, *Amer. Math. Soc.* 101, 1970.
- [10] F. John, Nonlinear Waves Equations, Formation of Singularities, *Pitcher Lectures in Math. Sciences*, Lehigh University, American Math. Society, 1990.
- [11] S.N. Kruzkov, First order quasilinear equations in several independent variables, *Mat. Sbornik (N.S.)*, 81 (1970), 228-255.
- [12] D.X. Kong, Q.Y. Sun and Y. Zhou, The equation for time-like extremal surfaces in Minkowski space $\mathbb{R}^{2+n}$, *J. Math. Phys.* 47 (2006), no. 1, 013503, 16 pages.
- [13] D.X. Kong and T. Yang, Asymptotic behavior of global classical solutions of quasilinear hyperbolic systems, *Comm. Part. Diff. Eqs.* 28 (2003), 1203-1220.
- [14] P.D. Lax, Hyperbolic systems of conservation laws II, *Comm. Pure Appl. Math.* 10 (1957), 537-566.
- [15] T.T. Li, Global Classical Solutions for Quasilinear Hyperbolic Systems, *Research in Appl. Math.* Vol. 32, Wiely/Masson, 1994.
- [16] T.T. Li, Y.J. Peng and J. Ruiz, Entropy solutions for linearly degenerate hyperbolic systems of rich type, *J. Math. Pures Appl.* 91 (2009), 553-568.
- [17] T.T. Li, Y. Zhou and D.X. Kong, Weak linear degeneracy and global classical solutions for general quasilinear hyperbolic systems, *Comm. Partial Diff. Equations*, 19 (1994), 1263-1317.
- [18] H. Lindblad, A remark on global existence for small initial data of the minimal surface equation in Minkowskian space time, *Proc. Am. Math. Soc.* 132 (2004), 1095-1102.
- [19] J.L. Liu and Y. Zhou, Asymptotic behavior of global classical solutions of diagonalizable quasilinear hyperbolic systems, *Math. Meth. Appl. Sci.* 30 (2007), 479-500.
- [20] T.P. Liu, Linear and nonlinear large-time behavior of solutions of general systems of hyperbolic conservation laws, *Comm. Pure Appl. Math.* 30 (1977), 767-796.

- [21] Y.J. Peng, Explicit solutions for 2×2 linearly degenerate systems, *Appl. Math. Letters*, 11 (1998), 75-78.
- [22] Y.J. Peng, Euler-Lagrange change of variables in conservation laws and applications, *Non-linearity*, 20 (2007), 1927-1953.
- [23] Y.J. Peng and Y.F. Yang, Well-posedness and long-time behavior of Lipschitz solutions to generalized extremal surface equations, *J. Math. Physics*, 52 (2011), 053702 (23 pages).
- [24] D. Serre, Richness and the classification of quasilinear hyperbolic systems, in :*Multidimensional Hyperbolic Problems and Computations*, Minneapolis, MN, 1989, in : *IMA Vol. Math. Appl.*, 29, Springer, New York, 1991, 315-333.
- [25] D. Serre, Systèmes de Lois de Conservation I-II, Diderot, Paris, 1996.
- [26] D. Serre, Hyperbolicity of the nonlinear models of Maxwell's equations, *Arch. Rat. Mech. Anal.* 172 (2004), 309-331.
- [27] B. Sévenec, Géométrie des systèmes hyperboliques de lois de conservation, *Mém. Soc. Math. France*, 56 (1994).
- [28] S.P.Tsarëv, On Poisson brackets and one-dimensional systems of hydrodynamic type, *Dokl. Akad. Nauk SSSR*, 282 (1985), 534-537.
- [29] S.P.Tsarëv, The geometry of Hamiltonian systems of hydrodynamic type. The generalized hodograph method (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 54 (1990), 1048-1068; translation in *Math. USSR-Izv.* 37 (1991), 397-419.
- [30] D.H. Wagner, Equivalence of the Euler and Lagrangian equations of gas dynamics for weak solutions, *J. Differential Equations*. 68 (1987), 118-136.